

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 495/04, A61K 31/38, 31/17 // (C07D 495/04, 333:00, 209:00) (C07D 495/04, 333:00, 221:00)	A1	(11) International Publication Number: WO 94/22871 (43) International Publication Date: 13 October 1994 (13.10.94)
(21) International Application Number: PCT/EP94/00917 (22) International Filing Date: 22 March 1994 (22.03.94) (30) Priority Data: 9306460.8 29 March 1993 (29.03.93) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex 9EP TW8 (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FORBES, Ian, Thomson [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldhar- bour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). MARTIN, Roger, Thomas [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). JONES, Graham, Elgin [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). (74) Agent: GIDDINGS, Peter, J.; Corporate Intellectual Property, SmithKline Beecham, Mundells, Welwyn Garden City, Hertfordshire AL7 1EY (GB).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: THIENO-INDOLE DERIVATIVES AS 5HT2c AND 5HT2b ANTAGONISTS (57) Abstract Novel thieno-indole compounds being 5HT2c and 5HT2b antagonists, processes for their preparation, compositions containing them and their use in the treatment of mammals.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

THIENO-INDOLE DERIVATIVES AS 5HT_{2c} AND 5HT_{2b} ANTAGONISTS


This invention relates to novel compounds having pharmacological activity, to processes for their preparation, to compositions containing them and to their use in the treatment of mammals.

P. Fludzinski et. al., J. Med. Chem. 1986 29 2415-2418 describes N-(1,2-dimethyl-3-ethyl-1H-indol-5-yl)-N'-(3-trifluoromethylphenyl)urea which shows selectivity for the rat stomach fundus serotonin receptor.

WO 92/05170 describes certain urea derivatives which are described as possessing 5HT_{1C} receptor antagonist activity. The 5HT_{1C} receptor has recently been reclassified as the 5HT_{2C} receptor [P. Hartig et al., Trends in Pharmacological Sciences (TIPS) 1993].

A structurally distinct class of compounds has now been discovered which have been found to have 5HT_{2C} receptor antagonist activity. Certain compounds of the invention also exhibit 5HT_{2B} antagonist activity. 5HT_{2C/2B} receptor antagonists are believed to be of potential use in the treatment of CNS disorders such as anxiety, depression, obsessive compulsive disorders, migraine, anorexia, Alzheimers disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus.

Accordingly, the present invention provides a compound of formula (I) or a salt thereof:



wherein:

P represents a quinoline or isoquinoline residue or a 5- or 6-membered aromatic heterocyclic ring containing up to three heteroatoms selected from nitrogen, oxygen or sulphur;

R₁ is hydrogen, C₁₋₆ alkyl, halogen, OR₇ or NR₈R₉ where R₇, R₈ and R₉ are

independently hydrogen or C₁₋₆ alkyl;

R₂ is hydrogen or C₁₋₆ alkyl;

R₃ is hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, or halogen;

R₄ is hydrogen or C₁₋₆ alkyl;

- 5 R₅ and R₆ are each independently hydrogen or C₁₋₆ alkyl; and
n is 2 or 3.

- 10 C₁₋₆alkyl groups, whether alone or as part of another group, may be straight chained or branched and are preferably C₁₋₃ alkyl, such as methyl, ethyl, *n*- and *iso*- propyl, most preferably methyl.

Preferably R₁ is hydrogen, C₁₋₆alkyl or halogen. Most preferably R₁ is hydrogen, methyl or bromo.

- 15 Preferably R₂ and R₃ are hydrogen.

Preferably R₅ and R₆ are both hydrogen and n is 2, that is to say, the group -(CR₅R₆)_n- forms an ethylene linkage. The group -(CR₅R₆)_n- can be attached to the 4- or 6-position of the benzothiophene ring.

20

The urea moiety can be attached to a carbon or, when available, a nitrogen atom of the ring P, preferably it is attached to a carbon atom.

- 25 Suitable moieties when the ring P is a 5- or 6-membered aromatic heterocyclic ring include pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, isothiazolyl, isoxazolyl, thiadiazolyl and triazolyl. When P is a quinoline or isoquinoline residue, the urea moiety can be attached at any position of the ring, preferably to the 4-position.

Preferably P is pyridyl, quinolyl or isothiazolyl.


30

The substituent R₃ can be attached to any vacant position in the phenyl part of the benzothiophene ring, that is to say, the 4-, 6- or 7-positions of the benzene ring. The substituent R₄ can be attached to the 2- or 3-positions of the thiophene ring.


- 35 Preferred compounds of formula (I) include;
7,8-Dihydro-6-(3-pyridylcarbamoyl)-thieno[3,2-e]-indole,

- 6,7-Dihydro-5-(3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 7,8-Dihydro-6-(5-quinolylcarbamoyl)-6H-thieno[3,2-e]indole,
 7,8-Dihydro-6-(3-Methyl-5-isothiazolylcarbamoyl)-6H-thieno[3,2-e]indole,
 6,7-Dihydro-5-(5-Bromo-3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 5 7,8-Dihydro-6-(2-methyl-4-quinolylcarbamoyl)-6H-thieno[3,2-e]indole,
 6,7-Dihydro-5-(2-methyl-4-quinolylcarbamoyl)-5H-thieno[2,3-f]indole,
 6,7-Dihydro-5-(4-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 and pharmaceutically acceptable salts thereof.
- 10 The compounds of the formula (I) can form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric, lactic, mandelic, tartaric and methanesulphonic.
- 15 Compounds of formula (I) may also form N-oxides, S-oxides or solvates such as hydrates, and the invention also extends to these forms. When referred to herein, it is understood that the term 'compound of formula (I)' also includes these forms.
- Certain compounds of formula (I) are capable of existing in stereoisomeric forms
 20 including enantiomers and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis. When R₂ is hydrogen or when R₁ is hydroxy or NR₇R₈ and at least one of R₇ and R₈ are hydrogen, the compounds of formula (I) may
 25 exist tautomerically in more than one form. The invention extends to all tautomeric forms of compounds of formula (I) and mixtures thereof.

The present invention also provides a process for the preparation of a compound of
 formula (I) or a pharmaceutically acceptable salt thereof, which process comprises
 30 (a) the coupling of a compound of formula (II);




35 with a compound of formula (III);



wherein A and B contain the appropriate functional group(s) necessary to form the moiety -NR₂'CO when coupled, wherein R₂' is R₂ as defined in formula (I) or a group convertible thereto, and the variables R₁', R₃', R₄', R₅' and R₆' are R₁, R₃, R₄, R₅ and R₆, respectively, as defined in formula (I), or groups convertible thereto, and thereafter optionally and as necessary in any appropriate order, converting any R₁', R₂', R₃', R₄', R₅' and R₆' when other than R₁, R₂, R₃, R₄, R₅ and R₆ to R₁, R₂, R₃, R₄, R₅ and R₆, interconverting R₁, R₂, R₃, R₄, R₅ and R₆, and forming a pharmaceutically acceptable salt, or

(b) cyclising a compound of formula (IV):



wherein R₁', R₂', R₃', R₅', and R₆' are as defined in formulae (II) and (III) and C and D contain the appropriate functional group(s) necessary to form the thiophene ring substituted by R₃' and R₄' as defined in formula (III), and thereafter optionally and as necessary in any appropriate order, converting any R₁', R₂', R₃', R₄', R₅' and R₆' when other than R₁, R₂, R₃, R₄, R₅ and R₆ to R₁, R₂, R₃, R₄, R₅ and R₆, interconverting R₁, R₂, R₃, R₄, R₅ and R₆, and forming a pharmaceutically acceptable salt.

Suitable examples of groups A and B include:

- (i) A is -N=C=O and B is -H,
- (ii) A is -NR₂'COL and B is -H,
- (iii) A is -NHR₂' and B is COL, or
- (iv) A is halogen and B is -CONHR₂'

wherein R₂' is as defined above and L is a leaving group. Examples of suitable leaving

groups L include imidazole, halogen such as chloro or bromo or phenoxy or phenylthio optionally substituted, for example with halogen.

5 When A is $-N=C=O$ and B is H the reaction is suitably carried out in an inert solvent for example dichloromethane or toluene at ambient temperature.

10 When A is $-NR_2'COL$ and B is H or when A is $-NHR_2'$ and B is COL, the reaction is suitably carried out in an inert solvent such as dichloromethane at ambient temperature optionally in the presence of a base, such as triethylamine or in dimethylformamide at ambient or elevated temperature.

When A is halogen and B is $CONHR_2'$, the reaction is suitably carried out in an inert solvent such as toluene at elevated temperature, optionally in the presence of a base.

15 Examples of compounds of formula (IV) which can be used in the preparation of benzothiophenes include those where C is $SCH_2CO_2R_5$ where R_5 is C_{1-6} alkyl and D is CHO, or C is $SCH_2CH(OR_5)_2$ and D is hydrogen.

20 Suitable examples of groups R_1' , R_3' and R_4' groups which are convertible to R_1 , R_3 and R_4 alkyl groups respectively, include acyl groups which are introduced conventionally and may be converted to the corresponding alkyl group by conventional reduction, such as using sodium borohydride in an inert solvent followed by hydrogenolysis in an inert solvent. Hydrogen substituents may be obtained from alkoxycarbonyl groups which may be converted to hydrogen by hydrolysis and decarboxylation. When R_1 is hydroxy it is
25 preferably protected in the compound of formula (II) as, for example, benzyloxy which is cleaved by hydrogenation. Suitable examples of a group R_2' which is convertible to R_2 include alkoxycarbonyl and benzyl or *para*-methoxybenzyl which are converted to R_2 is hydrogen using conventional conditions.

30 Interconversions of R_1 , R_2 , R_3 and R_4 are carried out by conventional procedures. For example, in the case where R_2 is hydrogen it is possible to introduce a C_{1-6} alkyl group at the R_2 position by conventional alkylation using 1 molar equivalent of a C_{1-6} alkyl halide and 1 molar equivalent of a suitable base in an inert solvent.

35 It should be appreciated that it may be necessary to protect any R_1 to R_4 hydrogen variables which are not required to be interconverted. Suitable protecting groups and

methods for their attachment and removal are conventional in the art of organic chemistry, such as those described in Greene T.W. 'Protective groups in organic synthesis' New York, Wiley (1981). It is preferable, however, to introduce and interconvert the groups R_1 to R_4 before coupling compounds of formulae (II) and (III) together, or cyclising the compound
 5 of formula (IV).

Compounds of formula (II) in which A is NHR_2' are known compounds or can be prepared analogously to known compounds, see, for example, WO 92/05170.

10 Compounds of formula (II) in which A is $-N=C=O$ may be prepared by treating a compound of formula (II) in which :

- i) A is amino, with phosgene or a phosgene equivalent, in the presence of excess base in an inert solvent.
- 15 ii) A is acylazide (i.e. CON_3), via the nitrene, by thermal rearrangement using conventional conditions (ref L.S. Trifonov et al, Helv. Chim. Acta 1987 70 262).
- iii) A is $CONH_2$, via the nitrene intermediate using conventional conditions.


Examples of phosgene equivalents include carbonyldiimidazole and phenyl chloroformate.
 20

Compounds of formula (II) in which A is $-NR_2'COL$ may be prepared by reacting a compound of formula (II) in which A is $-NHR_2'$ with phosgene or a phosgene equivalent, in an inert solvent, at low temperature, if necessary in the presence of one equivalent of a base such as triethylamine.

25 Compounds of formula (III) may be prepared:

(a) by cyclisation of compounds of formula (V), followed by reduction to the amine if necessary

30




wherein Q is CR^5R^6L , CR^5O or CO_2R where L is a leaving group and R^5 and R^6 are as

defined in formula (I), m is 1 or 2, R_3' , R_4' , R_5' , R_6' and B are as defined in relation to formula (III) above and R is aryl or alkyl group,

or (b) cyclisation of compounds of formula (VI)

5



wherein R_3' , R_5' , R_6' and B are as defined in relation to formula (III) above and C and D are as defined in relation to formula (IV) above.

10

The cyclisation of a compound of formula (V) may be suitably carried out in an inert solvent at ambient or elevated temperatures, optionally in the presence of a base. Reduction, if necessary, may be carried out using conventional reduction techniques. The cyclisation of a compound of formula (VI) may be suitably carried out using the procedures outlined for the cyclisation of a compound of formula (IV), above.

15

Compounds of formula (II) in which A is halogen and B is hydrogen are commercially available. Novel intermediates of formulae (III) and (IV) also form part of the invention.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative. N-oxides may be formed conventionally by reaction with hydrogen peroxide or percarboxylic acids.

Compounds of formula (I) and their pharmaceutically acceptable salts have 5HT_{2C/2B} receptor antagonist activity and are believed to be of potential use in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus.

30

Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders,

Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus.

- 5 The invention further provides a method of treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus, in mammals including humans, which
10 comprises administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

- In another aspect, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the
15 treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as hydrocephalus.

- 20 The present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

- 25 A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusable solutions or suspensions or suppositories. Orally administrable compositions
30 are generally preferred.

- Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well
35 known in normal pharmaceutical practice.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, 5 non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The 10 compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after 15 filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the 20 composition to facilitate uniform distribution of the compound.

The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

25 The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 20.0 mg, for example 0.2 to 5 mg; and such unit doses may be administered more than once a day, for example two or three a day, so that the total daily 30 dosage is in the range of about 0.01 to 100 mg/kg; and such therapy may extend for a number of weeks or months.

The following examples illustrate the invention.

Description 1**5-Nitrobenzo[b]thiophene (D1)**

5 Ethyl 5-nitrobenzo[b]thiophenecarboxylate was prepared and hydrolysed to the corresponding acid as described by S. Rossi and R. Trave (II Farmaco - Ed. Sci., 1960, 15, 396). 5-Nitrobenzo[b]thiophenecarboxylic acid (4.32 g, 19.4 mmol) was heated with copper powder (1.2 g, activated by heating for several hours at 160°C in vacuo) in quinoline (25 ml) at 180-190°C for 2h. After cooling, the mixture was diluted with ether
10 and washed thoroughly with 5N hydrochloric acid. The organic phase was dried and evaporated, and the crude product was recrystallised from ether to give the title compound (3.24 g, 77%), m.p. 142-145°C.

15 NMR (CDCl₃) δ: 7.52 (1H, d, J 6), 7.68 (1H, d, J 6), 8.00 (1H, d, J 8), 8.22 (1H, dd, J 8, 2), 8.74 (1H, d, J 2).

Description 2**4-Methyl-5-nitrobenzo[b]thiophene(D2)**

20 To a solution of 5-nitrobenzo[b]thiophene(D1) (1.79g, 10 mmol) in dry THF (100 ml), cooled to -20° C, was added methylmagnesium bromide (3M in ether, 6.66 ml, 20 mmol), such that internal temperature remained below -15° C. The mixture was then stirred at this temperature for 30 min. 2, 3-Dichloro-5,6-dicyanobenzoquinone (2.72 g, 12 mmol) was added and the mixture was stirred for 15 min. at room temperature. The mixture was then
25 poured into dilute acetic acid (aqueous) and extracted three times with dichloromethane. Extracts were combined, washed with sodium bicarbonate (twice) and water, dried and evaporated. The residue was chromatographed on silica gel (120g) eluted with 1:1 petrol/dichloromethane to give the title compound (1.35 g; 47% combined yield from two preparations), m.p. 104-105° C (ether/petrol).

30 NMR (CDCl₃)δ: 2.88 (3H, s), 7.60 (1H, d, J=6), 7.64 (1H, d, J=6), 7.81 (1H, d, J=10), 7.92 (1H, d, J=10)

Description 3

35 **4-(2-Hydroxyethyl)-5-nitrobenzo[b]thiophene (D3)**

A mixture of 4-methyl-5-nitrobenzothiophene (D2) (1.045 g, 5.5 mmol), paraformaldehyde (0.18 g, 6 mmol) and potassium hydroxide (0.5 M in ethanol, 1.2 ml, 0.6 mmol) in DMSO (10 ml) was stirred at room temperature for 4 h. The mixture was then diluted with ethyl acetate and washed thoroughly with water. The organic phase was dried and evaporated, and the residue was chromatographed on silica gel (60g) eluted with 0-1.5% methanol/dichloromethane, to give the title compound (0.92 g, 75%), m.p. 83-85°

NMR (CDCl₃)δ: 1.82 (1H, t, J=6), 3.51 (2H, t, J=6), 4.08 (2H, q, J=6), 7.66 (2H, s), 7.84 (1H, d, J=10), 7.91 (1H, d, J=10)

Description 4

4-(2-Methylsulphonyloxyethyl)-5-nitrobenzo[b]thiophene(D4)

A mixture of hydroxyethyl compound (D3) (1.03g, 4.6 mmol), methylsulphonyl chloride (0.4 ml, 5.2 mmol) and triethylamine (0.7 ml, 5 mmol) in dichloromethane (20 ml) was stirred for 1 h at room temperature. A further portion of triethylamine (0.2 ml) was added and stirring was continued for 30 min. Water (10 ml) was then added and the mixture was stirred for approx. 10 min before acidification with 5M hydrochloric acid and separation of phases. The organic phase was washed with dilute hydrochloric acid, dried and evaporated. The residue was recrystallised from ether to give the title compound (1.0 g, 72%), m.p. 108-110° C.

NMR (CDCl₃)δ: 2.97 (3H, s), 3.72 (2H, t, J=6), 4.67 (2H, t, J=6), 7.71 (1H, d, J=6), 7.74 (1H, d, J=6), 7.92 (1H, d, J=10), 7.99 (1H, d, J=10)

Description 5

7,8-Dihydrothieno[3,2-e]indole (D5)

A mixture of the methanesulphonate (D4) (1.0g, 3.3mmol) and 10% palladium on charcoal (0.1 g) in ethanol (50 ml) was stirred under hydrogen at 47 psi for 6.5 h. A further 50 mg of palladium/charcoal was added and hydrogenation continued for 7 h. The mixture was filtered through kieselguhr and evaporated. The residue was partitioned between dichloromethane and water and the organic phase was extracted with dilute hydrochloric acid. The combined aqueous extracts were basified with 10% sodium hydroxide and extracted with DCM. The organic extract was dried and evaporated to give the title compound (0.18 g, 31%) as a greenish oil.

NMR (CDCl₃) δ : 3.31 (2H, t, J=8), 3.78 (2H, t, J=8), 4.75 (broad s), 6.96 (1H, d, J=10), 7.18 (1H, d, J=6), 7.48 (1H, d, J=6), 7.60 (1H, d, J=10)

5 Description 6

*2-Bromo-4-methylbenzenecarboxaldehyde (D6)

Sodium nitrite (45.0g, 650 mmoles) in water (80 ml) was added dropwise to a mixture of 2-bromo-4-methylaniline (100.0g, 530 mmoles), concentrated hydrochloric acid (125 ml) and ice/water (400 ml) over 30 mins keeping the temperature <5° C. The mixture was stirred for a further 30 mins at 0° C and sodium acetate trihydrate (50.0g, 360 mmoles) was added.

The diazonium solution/suspension was then added portionwise via a transfer tube to a solution of formaldoxime [formaldoxime prepared from paraformaldehyde (32.5g, 1000 mmoles) and hydroxylamine hydrochloride (74.0g, 1050 mmoles) by heating under reflux in water (400 ml) with sodium acetate trihydrate (150.0g, 1080 mmoles), copper sulphate pentahydrate (16.3g), sodium sulphite (2.2g) and anhydrous sodium acetate (210g, 2560 mmoles) for 20 mins] with vigorous stirring keeping the temperature <15° C (Care! reaction froths greatly). The mixture was stirred at ambient temperature for 1 hr and concentrated hydrochloric acid (470ml) was added. The mixture was then heated under reflux for 1 hr.

The product was steam distilled from the reaction for 3 hr and then extracted into diethyl ether (2x1000 ml), washed with saturated aqueous sodium bicarbonate solution, dried (Na₂SO₄) and evaporated to dryness. Vacuum distillation at 5 mm Hg gave the title compound (57g, 53%) b.p. 102-108° C

NMR (CDCl₃) δ : 2.41 (3H, s), 7.22 (1H, d, J 8), 7.48 (1H, s), 7.81 (1H, d, J 8), 10.30 (1H, s).

* S.D. Jolad, S. Rajagopal, Org Synth Coll Vol, V, P.139, (1973)

Description 7

35 2-Bromo-4-methyl-5-nitrobenzenecarboxaldehyde (D7)

Concentrated nitric acid (36ml, 570 mmoles) was added dropwise to 2-bromo-4-methylbenzene-carboxaldehyde (D6) (57g, 285 mmoles) in concentrated sulphuric acid (300 ml) with stirring keeping the temperature <15° C. The mixture was stirred for a further 20 mins then poured onto ice/water (250 ml) with vigorous stirring. The mixture
5 was extracted with 5% methanol/chloroform (3 x 1000 ml), washed with saturated sodium bicarbonate solution, filtered and evaporated to dryness. The residue was recrystallised from ethyl acetate/60-80 petrol to give the title compound (44.2g, 63%).

NMR (CDCl₃) δ : 2.63 (3H, s), 7.68 (1H, s), 8.49 (1H, s), 10.30 (1H, s).

10

Description 8

Ethyl-5-nitro-6-methylbenzo[b]thiophene-2-carboxylate (D8)

Ethyl-2-mercaptoacetate (20 ml, 182 mmoles) was added to a solution of sodium ethoxide
15 [prepared by treating ethanol (150 ml) with sodium metal (4.3g, 180 mmoles)] keeping the temperature <5° C. After 20 mins ethanol (300 ml) was added followed by 2-bromo-4-methyl-5-nitrobenzene-carboxaldehyde (D7) (44.2g, 180 mmoles) portionwise over 10 mins resulting in a dense yellow precipitate. More ethanol (200 ml) was added and the mixture heated under reflux for 3 hr. The mixture was allowed to cool and the ethanol
20 was removed *in vacuo*. The residue was partitioned between dichloromethane and water, separated, dried (Na₂SO₄) and evaporated to dryness to give the title compound (46.0g, 96%).

NMR (CDCl₃) δ : 1.42 (3H, t, J 6), 2.72 (3H, s), 4.42 (2H, q, J 6), 7.80 (1H, s), 8.10 (1H,
25 s), 8.53 (1H, s).

Description 9

5-Nitro-6-methylbenzo[b]thiophene (D9)

Ethyl-5-nitro-6-methylbenzo[b]thiophene-2-carboxylate (D8) (46.0g, 170 mmoles) in ethanol (500 ml) and water (300 ml) was heated under reflux with sodium hydroxide (27.0 g, 680 mmoles) for 3 hours. The mixture was then allowed to cool and most of the ethanol was removed *in vacuo*. The aqueous was then poured into a mixture of dilute hydrochloric acid (250 ml) and ice/water (100 ml) with vigorous stirring. The resulting
35 fine precipitate was filtered off and dried. The dry solid was then heated in quinoline (300 ml) with copper powder (25 g, 390 mmoles) to 190° C for 3 hrs with stirring. The mixture

was then cooled and diluted with diethyl ether (100 ml), filtered through kieselguhr. The filtrate was washed with dilute hydrochloric acid (3 x 500 ml), water (500 ml) and saturated sodium chloride (500 ml), separated, dried (Na₂SO₄), filtered and evaporated to dryness to give the title compound (29.6g, 88%)

5

NMR (CDCl₃) δ : 2.70 (3H, s), 7.40 (1H, d, J 6), 7.55 (1H, d, J 6), 7.80 (1H, s), 8.49 (1H, s).

Description 10

10 2-(5-Nitro-6-benzo[b]thienyl)acetaldehyde (D10)

5-Nitro-6-methylbenzo[b]thiophene (D9) (29.6g, 153 mmoles) in dimethyl formamide (300 ml) was treated with dimethylformamide dimethyl acetal (61 ml, 460 mmoles) and pyrrolidine (25.7 ml, 300 mmoles) and heated to 150° C for 3 hrs under argon. The mixture was then cooled and evaporated to dryness. Toluene (600 ml), water (800 ml) and dilute hydrochloric acid (300 ml) were added to the residue and the mixture heated under reflux for 30 mins. The mixture was cooled and extracted with ethyl acetate (2 x 600 ml). The organic layer was washed with water (500 ml), dried (Na₂SO₄) and evaporated to dryness. Flash chromatography on TLC silica gel eluting with dichloromethane gave the title compound (26.4g, 78%)

15
20

NMR (CDCl₃) δ : 4.21 (2H, s), 7.48 (1H, d, J 6), 7.63 (1H, d, J 6), 7.79 (1H, s), 8.63 (1H, s), 9.90 (1H, s)

25 Description 11

2-(5-Nitro-6-benzo[b]thienyl)ethanol (D11)

2-(5-Nitro-6-benzo[b]thienyl)acetaldehyde (D10) (25.3g, 114 mmoles) in ethanol was treated with sodium borohydride (8.7g, 228 mmoles) with stirring at ambient temperature. Stirring was continued for 1 hr then most of the ethanol was removed *in vacuo*. Water (500 ml) was added. The mixture was then acidified by careful addition of dilute hydrochloric acid (gas evolution) and then extracted with ethyl acetate (2 x 400 ml). The combined organics were washed with water (500 ml) and saturated sodium chloride solution (500 ml), separated, dried (Na₂SO₄) and evaporated to dryness. This gave the title compound (25.0g, 98%).

30
35

NMR (CDCl₃) δ : 3.28 (2H, t, J 8), 3.95 (2H, t, J 8), 7.40 (1H, d, J 6), 7.58 (1H, d, J 6), 7.89 (1H, s), 8.45 (1H, s).

Description 12

5 2-(5-Nitro-6-benzo[b]thienyl)ethyl methane sulphonate (D12)

2-(5-Nitro-6-benzo[b]thienyl)ethanol (D11) (25.0 g, 112 mmoles) in dichloromethane (500 ml) was treated with triethylamine (16.4 ml, 118 mmoles). Methanesulphonyl chloride (9.2 ml, 118 mmoles) was then added dropwise with stirring over 15 mins. The
10 mixture was stirred for a further 1 hr then washed with saturated aqueous sodium bicarbonate (2 x 500 ml), dried (Na₂SO₄) and evaporated to dryness to give the title compound (19.6g, 88%).

NMR (CDCl₃) δ : 2.98 (3H, s), 3.45 (2H, d, J 8), 4.60 (2H, d, J 8), 7.45 (1H, d, J 6), 7.63
15 (1H, d, J 6), 7.89 (1H, s), 8.51 (1H, s)

Description 13

6,7-Dihydro-5H-thieno[2,3-f]indole (D13)

20 The methanesulphonate (D12) (24.0g, 80 mmoles) was hydrogenated at atmospheric pressure and at 50° C for 3 hrs in ethyl acetate (600 ml) with triethylamine (40 ml) over a 10% palladium on charcoal catalyst. The catalyst was then filtered off and the filtrate washed with saturated aqueous sodium bicarbonate, dried (Na₂SO₄) and evaporated to dryness to give the title compound as an off-white solid (12.7g, 91%)

25 NMR (CDCl₃) δ : 3.12 (2H, t, J 8), 3.40-3.60 (1H, br s), 3.65 (2H, t, J 8), 7.01 (1H, s), 7.11 (1H, d, J 6), 7.25 (1H, d, J 6), 7.53 (1H, s)

Example 1

30 7,8-Dihydro-6-(3-pyridylcarbamoyl)-thieno[3,2-e]-indole (E1)

A solution of nicotinic acid azide (0.16 g, 1.1 mmol) in toluene (7 ml) was heated under reflux for 1.5 h. The solution was cooled and thienoindole (D5) (0.18g, 1.03 mmol) was added in dichloromethane (7 ml) containing a little dimethylformamide. The mixture was
35 stirred at room temperature for 2.5 h. The mixture was then washed with water, and the aqueous phase was extracted with dichloromethane/methanol. The toluene phase and the dichloromethane extract were combined, dried and evaporated. The residue was

recrystallised from ethanol/water to give the title compound (0.18 g, 59%) m.p. 192-197° C.

Found: C, 64.97; H, 4.63; N, 14.05%

5 $C_{16}H_{13}N_3OS$ requires: C, 65.06; H, 4.44; N, 14.23%

NMR(D_6 -DMSO) δ : 3.45(2H, t, J=8), 4.30 (2H, t, J=8), 7.33 (1H, d, J=6), 7.36 (1H, d, J=6), 7.78 (1H, d, J=6), 7.80 (1H, d, J=6), 8.01 (1H, m, J=7), 8.07 (1H, d, J=7), 8.24 (1H, d, J=4), 8.74 (1H, s), 8.78 (1H, s)

10

Example 2

6,7-Dihydro-5-(3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole (E2)

15 Nicotiny azide (0.06g, 0.4 mmol) in toluene (3ml) was heated under reflux for 1 h, then cooled. 6,7-Dihydro-5H-thieno[2,3-f]indole (0.07g, 0.4 mmol) in dichloro- methane was added and the mixture was stirred for 1 h at room temperature. A little petrol was added and the solid product was filtered off, washed with petrol and dried *in vacuo* to give the title compound (0.085g, 72%), m.p. 183-185° C.

20 Found: C, 64.94; H, 4.73; N, 14.26 %

$C_{16}H_{13}N_3OS$ requires C, 65.06; H, 4.44; N, 14.23%

NMR (D_6 -DMSO) d: 3.28 (2H, t, J=9), 4.22 (2H, t, J=9), 7.34 (1H, m), 7.37 (1H, d, J=5), 7.63 (1H, d, J=5), 7.77 (1H, s), 8.01 (1H, d, J=8), 8.23 (1H, d, J=5), 8.33 (1H, s), 8.75 (2H, m).

25

Example 3

7,8-Dihydro-6-(5-quinolylcarbamoyl)-6H-thieno[3,2-e]indole (E3)

30 5-Aminoquinoline (0.27g, 1.9 mmol) and 1,1'-carbonyldiimidazole (0.31g, 1.9 mmol) were stirred at reflux under Ar in dry dichloromethane (10 ml) for 2.5h. The mixture was evaporated to dryness, and the residue was dissolved in dry DMF (10 ml). 7,8-Dihydro-6H-thieno[3,2-e]indole (D5) (0.30g, 1.7 mmol) was added, and this mixture was stirred at 115° C for 1h, when it was poured into water (200 ml). The precipitate was filtered off
35 and dried *in vacuo*. Chromatography on silica gel, eluting with 0-1% methanol/ethyl acetate, gave the title compound (0.31g, 52%) as a greenish-grey powder, m.p. 243-4° C.

NMR (DMSO-d₆) (δ): 3.49 (2H, t, J 8), 4.44 (2H, t, J 8), 7.36 (1H, d, J 5), 7.53 (1H, dd, J 8, 4), 7.64 (1H, d, J 7), 7.7-7.85 (3H, m), 7.92 (1H, d, J 8), 8.03 (1H, d, J 9), 8.48 (1H, d, J 8), 8.85 (1H, s), 8.92 (1H, m)

- 5 Found: C, 66.7; H, 4.4; N, 11.7%
C₂₀H₁₅N₃OS · 0.75H₂O requires C, 66.9; H, 4.6; N, 11.7%

Example 4

7,8-Dihydro-6-(3-Methyl-5-isothiazolylcarbamoyl)-6H-thieno[3,2-e]indole (E4)

- 10 This was prepared from 5-amino-3-methylisothiazole hydrochloride (0.26g, 1.7mmol) and 7,8-dihydro-6H-thieno[3,2-e]indole (D5), following the procedure of Example 3, but including 1 equivalent of triethylamine in the initial step. Recrystallisation from methanol/dichloromethane gave the desired material.

- 15 NMR (DMSO-d₆) (δ): 2.30 (3H, s), 3.47 (2H, t, J 8), 4.26 (2H, t, J 8), 6.77 (1H, s), 7.36 (1H, d, J 6), 7.8 (2H, m), 8.06 (1H, d, J 8), 10.55 (1H, b s).

Found: C, 57.1; H, 4.3; N, 13.4%. C₁₅H₁₃NOS₂ requires C, 57.1; H, 4.2; N, 13.3%

20 **Example 5**

6,7-Dihydro-5-(5-Bromo-3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole (E5)

This was prepared from 6,7-dihydro-5H-thieno[2,3-f]indole (D) (0.28g, 1.6 mmol) and 5-bromonicotinoyl azide (0.38g, 1.7 mmol), following the procedure of Example 2.

- 25 Recrystallisation of the crude product from ethanol/petroleum ether (b.p. 60-80°C) gave the title compound (0.16g, 34%) as an off-white solid, m.p. 203-5° C decomp.

NMR (DMSO-d₆) (δ): 3.34 (2H, t, J 8), 4.23 (2H, t, J 8), 7.39 (1H, d, J 4), 7.65 (1H, d, J 4), 7.80 (1H, s), 8.35 (3H, m), 8.78 (1H, d, J 2), 8.97 (1H, s)

30

Found: C, 51.2; H, 3.4; N, 11.0%. C₁₆H₁₂BrN₃OS requires: C, 51.4; H, 3.2; N, 11.2%

Example 6

7,8-Dihydro-6-(2-methyl-4-quinolylcarbamoyl)-6H-thieno[3,2-e]indole (E6)

35

To an ice-cooled solution of 1,1'-carbonyl diimidazole (0.47g, 2.86 mmol) in dry

dichloromethane (15 ml) was added 4-aminoquinoline (0.41g, 2.6 mmol) in dichloromethane (30 ml). The mixture was stirred for 1.75 h while warming from 0°C to room temperature. Solvent was then evaporated and replaced by dry DMF (10 ml). A solution of 7,8-dihydro-6H-thieno[3,2-e]indole (D5, 0.45g, 2.57 mmol) in dry DMF (10 ml) was added, and the mixture was stirred for 1.25h at 100-120° C. The mixture was then cooled and poured into water. The solid product was filtered off, washed with water and dried, then recrystallised twice from dichloromethane/petrol to give the title compound (0.35g, 38%), m.p. 229-231° C.

10 NMR (DMSO-d₆) (δ): 2.65 (3H, s), 3.49 (2H, t, J 8), 4.50 (2H, t, J 8), 7.38 (1H, d, J 6), 7.54 (1H, t, J 7), 7.72 (1H, t, J 7), 7.80 (3H, m), 7.92 (1H, d, J 7), 8.07 (1H, d, J 7), 8.18 (1H, d, J 7), 8.87 (1H, s)

Found: C, 70.04; H, 5.02; N, 11.66%

15 C₂₁H₁₇N₃OS requires C, 70.17; H, 4.77; N, 11.69%

Example 7

6,7-Dihydro-5-(2-methyl-4-quinolylcarbamoyl)-5H-thieno[2,3-f]indole (E7)

20 The title compound was prepared by the method of Example 3 from 2-methyl-4-aminoquinoline (0.46g, 29 mmoles) and 6,7-dihydro-5H-thieno [2,3-f] indole (D13) (0.5g, 29 mmoles). Recrystallisation from methanol/dichloromethane gave the title compound as white crystals (0.51g, 49%) m.p. 237-8° C.

25 NMR (DMSO-d₆) δ: 2.60 (3H, s), 3.30 (2H, t, J 8), 4.40 (2H, t, J 8), 7.35 (1H, d, J 6), 7.45-7.55 (1H, m), 7.60-7.95 (5H, m), 8.10-8.21 (1H, m), 8.35 (1H, s), 8.30-8.45 (1H, brs).

Found: C, 69.87; H, 4.89; N, 11.70%

30 C₂₁H₁₇N₃OS requires: C, 70.17; H, 4.77; N, 11.69%

Found M⁺, C₂₁H₁₇N₃OS requires 359

Example 8

6,7-Dihydro-5-(4-pyridylcarbamoyl)-5H-thieno[2,3-f]indole (E8)

35

The title compound was prepared by the method of Example 3 from 4-aminopyridine

(0.27g, 29 mmoles) and 6,7-dihydro-5H-thieno[2,3-f]indole (0.5g, 29 mmoles).

Recrystallisation from methanol/dichloromethane gave the title compound as light green crystals (0.45g, 53%) m.p. >240° C.

- 5 NMR (DMSO-d₆) δ : 3.30 (2H, t, J 8), 4.22 (2H, t, J 8), 7.39 (1H, d, J 6), 7.59-7.68 (3H, m), 7.80 (1H, s), 8.30-8.44 (3H, m), 8.95 (1H, s).

Found: C, 63.91; H, 4.63; N, 14.10%

C₁₆H₁₃N₃OS¹/₃H₂O requires: C, 63.79; H, 4.54; N, 13.95%

- 10 Found M⁺ 295, C₁₆H₁₃N₃OS requires 295

Pharmacological Data

Reversal of MCPP-induced Hypolocomotion

- 5 Administration of m-(chlorophenyl)piperazine (mCPP) to rats induces hypolocomotion (Kennett and Curzon 1988, Luckie *et al.* 1989) as seen with the related drug 1-(m-trifluoromethylphenyl)piperazine (TFMPP) (Lucki and Frazer 1982, Kennett and Curzon 1988). This effect was blocked by the non specific 5-HT_{2C}/5-HT_{2A} receptor antagonists mianserin, cyproheptadine and metergoline and perhaps by mesulergine. It was not
10 blocked by the 5-HT_{2A} receptor antagonists ketanserin and ritanserin at relevant doses (Kennett and Curzon 1991) nor by antagonists of 5-HT_{1A}, 5-HT_{1B}, 5-HT₃, α_2 adrenoceptors or dopamine D₂ receptors. The effect of mCPP is therefore considered to be mediated by 5-HT_{2C} receptors (Kennett and Curzon 1988) as confirmed by subsequent studies (Lucki *et al.*, 1989). Since mCPP causes hypolocomotion when infused into the
15 cerebral ventricles this effect is probably centrally mediated (Kennett and Curzon 1988).

- mCPP-induced hypolocomotion was measured in automated locomotion cages of dimensions 56 cm long x 16½ cm wide x 25 cm high and made of black perspex. Two photobeams traversed the width of the cages at either end at ground level. Sequential
20 breaking of these beams allowed the measurement of cage transits.

- Male Sprague Dawley rats (200-250g) (Charles River) were housed in groups of six. They were given drugs orally 1h pretest and 40 mins later mCPP (7 mg/kg i.p.). After a further 20 min they were placed in individual automated cages in groups of four under red light in
25 an adjacent room. After 10 min the test was terminated. Reversal of mCPP-induced hypolocomotion was considered as evidence of *in vivo* central 5-HT_{2C} receptor antagonist properties.

- Kennett, G.A., Curzon, G., (1988). *Brit. J. Pharmacol.* **94**, 137-147.
30 Kennet G.A., Curzon, G., (1991). *Brit.J. Pharmacol.* **103**, 2016-2020.
Lucki, I., Frazer, A., (1982) *Am. Soc. Neurosci.* 8(abstr.), 101.
Lucki, I., Ward, M.R., Frazer, A., (1989). *J.Pharmacol. Exp. Therap.* **249**, 155-164.

- The compounds of examples 1, 2 and 5 had ID50 values of 1.4 to 9.6 mg/kg p.o.
35

[³H]-mesulergine binding to rat or human 5-HT_{2C} clones expressed in 293 cells in vitro

- 5 Evidence from the literature suggests that 5-HT_{2C} antagonists may have a number of therapeutic indications including the treatment of anxiety, migraine, depression, feeding disorders and obsessive compulsion disorders. (Curzon and Kennett, 1990; Fozard and Gray, 1989) and Alzheimer's Disease (Lawlor, 1989, J. Arch. Gen. Psychiat. Vol. 46 p.542).

10

The affinity of test drugs for the 5-HT_{2C} binding site can be determined by assessing their ability to displace [³H]-mesulergine from 5-HT_{2C} clones expressed in 293 cells (Julius *et al.*, 1988). The method employed was similar to that of Pazos *et al.*, 1984.

- 15 The cells suspension (50ml) was incubated with [³H]-mesulergine (0.5nM) in Tris HCl buffer (pH 7.4) at 37°C for 30 minutes. Non-specific binding was measured in the presence of mianserin (10⁻⁶M). Ten concentrations of test drug (3 x 10⁻⁹ to 10⁻⁴M final concentration) were added in a volume of 50ml. The total assay volume was 500ml. Incubation was stopped by rapid filtration using a Brandel cell harvester and
- 20 radioactivity measured by scintillation counting. The IC₅₀ values were determined using a four parameter logistic program (DeLean 1978) and the pK_i (the negative logarithm of the inhibition constant) calculated from the Cheng Prusoff equation where:

25

$$K_i = \frac{IC_{50}}{1 + \frac{C}{K_d}}$$

- 30 K_i = inhibition constant.
C = concentration of [³H]-mesulergine
K_d = Affinity of mesulergine for 5-HT_{2C} binding sites.

Curzon, G.A. and Kennett, G.A. (1990). TIPS, Vol. 11, 181-182.

- 35 Fozard, J.R. and Gray, J.A. (1989). TIPS, Vol. 10, 307-309.
Pazos, A. *et al.* (1984). Eur. J. Pharmacol., 106, 531-538.

Julius *et al.* (1988) Science 241, 558-564

DeLean A, Munson P.J., Rodbaud D (1978) Am. J. Physiol 235, E97-E102.

The compounds of examples 1 to 8 had pK_i values from 5.1 to 8.4.

5

Geller-Seifter Procedure

Potential anxiolytic properties are evaluated using the Geller-Seifter procedure based on that originally described by Geller and Seifter, (1960) Psychopharmacologia, 1, 482-492.

10 This procedure has been shown to be selective for drugs with anxiolytic properties (Cook and Sepinwall, (1975) "Mechanism of Action of Benzodiazepines" ed. Costa, E. and Greengard, P., Raven Press, New York, pp. 1-28).

15 Rats are trained on a variable interval 30 sec schedule (VI30) to press a lever in order to obtain food reward. The 5 min sessions of the VI30 schedule alternate with 2-5 min of a schedule (FR5) in which every 5th lever press is followed by presentation of a food pellet paired with a 0.5 sec mild footshock. The total study lasts approximately 30 mins. Rats typically respond with high rates of lever pressing under the VI30 schedule and low response rates under the FR5 'conflict' session. Anxiolytic drugs increase the suppressed
20 response rates of rats in a 'conflict' session.

Drugs are administered intraperitoneally or orally to groups of 3-8 rats 30 min before testing. The results are expressed as the percentage increase in the square root of the total number of lever presses in the FR5 'conflict' session. Square root transformation is
25 necessary to normalise the data for statistical analysis using parametric methods.

The compounds of Examples 1 and 2 showed a significant increase in responding in the 'conflict' session at dose levels in the range 1 to 2 mg/kg p.o.

30 Rat Fundus: 5-HT_{2B} Receptors

The 5-HT receptor in the rat fundic strip (RFS) has been characterised as 5-HT_{2B}. Hence this tissue may be used to assess the 5-HT_{2B} antagonist properties of compounds.


35 Whole stomachs were obtained from male CD Rats (Charles River, 250-350g). Strips of fundus (2cm x 0.5 cm) were cut from the greater curvature and the mucosae carefully

removed. Tissues were then further dissected into smaller strips (2mm x 20mm) which were mounted in organ baths containing oxygenated Tyrodes solution at 37° C containing indomethacin (3 μ M). Preparations were maintained under a resting tension of 0.5g and exposed to the irreversible MAO inhibitor pargyline (100 μ M for 30 minutes followed by washout). Over a 1h equilibration period, rat fundic strips were challenged with 1×10^{-8} M 5-HT at 15 minute intervals until cumulative concentration-effect curve to the standard agonist 5-HT (1×10^{-10} upwards) was constructed to determine the individual sensitivity of each preparation. A further concentration-effect curve to either 5-HT, or other agonists was constructed no sooner than 1h after completion of the previous curve. When necessary tissues were equilibrated with the antagonists over this one hour period. Antagonists affinities are expressed as pA_2 estimates.

The compound of Example 2 has a pA_2 value of 7.97.

Claims:

1. A compound of formula (I) or a salt thereof:



wherein:

P represents a quinoline or isoquinoline residue or a 5- or 6-membered aromatic heterocyclic ring containing up to three heteroatoms selected from nitrogen, oxygen or sulphur;

R₁ is hydrogen, C₁₋₆ alkyl, halogen, OR₇ or NR₈R₉ where R₇, R₈ and R₉ are independently hydrogen or C₁₋₆ alkyl;

R₂ is hydrogen or C₁₋₆ alkyl;

R₃ is hydrogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, or halogen;

R₄ is hydrogen or C₁₋₆ alkyl;

R₅ and R₆ are each independently hydrogen or C₁₋₆ alkyl; and

n is 2 or 3.

2. A compound according to claim 1 in which R₁ is hydrogen, C₁₋₆alkyl or halogen.

3. A compound according to claim 1 or 2 in which R₂ and R₃ are hydrogen.

4. A compound according to any one of claims 1 to 3 in which R₅ and R₆ are both hydrogen and n is 2.

5. A compound according to any one of claims 1 to 4 in which P is pyridyl, quinolyl or isothiazolyl.

6. A compound according to claim 1 which is:

7,8-Dihydro-6-(3-pyridylcarbamoyl)-thieno[3,2-e]-indole,
 6,7-Dihydro-5-(3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 7,8-Dihydro-6-(5-quinolylcarbamoyl)-6H-thieno[3,2-e]indole,
 7,8-Dihydro-6-(3-Methyl-5-isothiazolylcarbamoyl)-6H-thieno[3,2-e]indole,
 5 6,7-Dihydro-5-(5-Bromo-3-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 7,8-Dihydro-6-(2-methyl-4-quinolylcarbamoyl)-6H-thieno[3,2-e]indole,
 6,7-Dihydro-5-(2-methyl-4-quinolylcarbamoyl)-5H-thieno[2,3-f]indole,
 6,7-Dihydro-5-(4-pyridylcarbamoyl)-5H-thieno[2,3-f]indole,
 and pharmaceutically acceptable salts thereof.

10


7. A compound according to any one of claims 1 to 6 for use in therapy.

8. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 6 and a pharmaceutically acceptable carrier or excipient.

15


9. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises:

(a) the coupling of a compound of formula (II);



20

with a compound of formula (III);



25


wherein A and B contain the appropriate functional group(s) necessary to form the moiety -NR₂'CO when coupled, wherein R₂' is R₂ as defined in formula (I) or a group convertible thereto, and the variables R₁', R₃', R₄', R₅' and R₆' are R₁, R₃, R₄, R₅ and R₆, respectively, as defined in formula (I), or groups convertible thereto, and thereafter optionally and as necessary in any appropriate order, converting any R₁', R₂', R₃', R₄', R₅' and R₆' when other than R₁, R₂, R₃, R₄, R₅ and R₆ to R₁, R₂, R₃, R₄, R₅ and R₆.

30

interconverting R_1 , R_2 , R_3 , R_4 , R_5 and R_6 , and forming a pharmaceutically acceptable salt, or

(b) cyclising a compound of formula (IV):

5



wherein R_1' , R_2' , R_3' , R_5' , and R_6' are as defined in formulae (II) and (III) and C and D contain the appropriate functional group(s) necessary to form the thiophene ring substituted by R_3' and R_4' as defined in formula (III), and thereafter optionally and as necessary in any appropriate order, converting any R_1' , R_2' , R_3' , R_4' , R_5' and R_6' when other than R_1 , R_2 , R_3 , R_4 , R_5 and R_6 to R_1 , R_2 , R_3 , R_4 , R_5 and R_6 , interconverting R_1 , R_2 , R_3 , R_4 , R_5 and R_6 , and forming a pharmaceutically acceptable salt.

10

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 94/00917

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D495/04 A61K31/38 A61K31/17 //(C07D495/04, 333:00,
209:00), (C07D495/04, 333:00, 221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,92 05170 (SMITHKLINE BEECHAM PHARMACEUTICALS) 2 April 1992 cited in the application ---	1-9
P,A	WO,A,93 18028 (SMITHKLINE BEECHAM PHARMACEUTICALS) 16 September 1993 ---	1-9
P,X	WO,A,94 04533 (SMITHKLINE BEECHAM PHARMACEUTICALS) 3 March 1994 ---	1-9
A	J.MED.CHEM. vol. 29 , 1986 pages 2415 - 2418 FLUDZINSKI, P. ET AL. '2.3-Dialkyl(dimethylamino)indoles: Interaction with...' cited in the application -----	1-9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

26 July 1994

Date of mailing of the international search report

09.08.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Stellmach, J

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 94/00917

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9205170	02-04-92	AU-B- 642041	07-10-93
		AU-A- 8503891	15-04-92
		CA-A- 2091246	14-03-92
		EP-A- 0550507	14-07-93
		JP-T- 6500551	20-01-94
		US-A- 5328922	12-07-94
WO-A-9318028	16-09-93	NONE	
WO-A-9404533	03-03-94	AU-B- 4704693	15-03-94